

CROP	DISEASE	RATE	WITHHOLDING PERIOD	WHEN TO APPLY
Grapes	Grey Mould (Botrytis)	150-200 ml/100 litres	1 day	Apply once or twice in the period from veraison to harvest using alternative, non dicarboximides for the rest of the season.
Stonefruit	Brown Rot (Monilinia spp)	150 ml/100 litres	1 day	Apply 2-3 sprays at 7-10 day intervals (more frequently following heavy rainfall) in the four week period prior to harvest. Use alternative non dicarboximide fungicides for the rest of the season.
Kiwifruit	Botrytis & Sclerotinia Fruit Rots	150ml/100 litres	1 day (24 hrs)	Sclerotinia control: Apply at flowering and again in January if required. Botrytis control: To reduce losses from Botrytis storage rot, apply one day before harvest. NB. Applications at flowering for Sclerotinia control will also suppress Botrytis. DO NOT make more than two applications in any one season.
Tangelos	Alternaria leaf spot (Alternaria citri)	200ml/100 litres	14 days	To reduce losses from Alternaria citri; Apply when last seasons crop has been harvested completely. Apply in a 21 28 day programme with alternative products from the start of flowering. Use the shorter interval during periods of high disease pressure.
Strawberries & other Berryfruit **	Botrytis	200ml/100 litres	1 day	Apply up to twice per season in a programme with alternative products on a 7-10 day schedule from the start of flowering. Reserve use for when infection pressure is high particularly in the pre harvest period.
Asparagus	Stemphylium	3 litres/hectare		Apply between close up and the end of February at first appearance of the disease on a 21-28 day schedule. Use the shorter interval during periods of prolonged wet weather.

** Do not use on Rabbiteye blueberry cvs.

Experience has shown Rovral Flo to be well tolerated by target field crops but selectivity should be checked before use on new cultivars or introductions

It is an offence for users of this product to cause residues exceeding the relevant MRL in the New Zealand (Maximum Residues Limits of Agricultural Compounds) Food Standards.

Rovral Flo is a contact fungicide of the dicarboximide group.

It is recommended as a foliar spray for the control of:

- Botrytis - Grey mould - in grapes, berryfruit and kiwifruit.
- Alternaria leaf spot - in tangelos.
- Monilinia - Brown rot - in stonefruit.
- Sclerotinia - Field rot - in kiwifruit.
- Stemphylium - Needle blight - in asparagus.

NOTE: Resistance to this dicarboximide fungicide and related products could develop from excessive use. To minimise this risk use strictly in accordance with label instructions.

It is recommended that when used for Botrytis control no more than two applications of Rovral Flo are made per season and that alternative non dicarboximide products are used at other key times in the programme.

Do not use this product if resistance monitoring tests show a high level of tolerant strains are present before application is planned.

Use Rovral Gold on glasshouse crops.

IT IS AN OFFENCE TO USE THIS PRODUCT ON ANIMALS.

DIRECTIONS FOR USE

Three-quarter fill the spray tank, add the required quantity of Rovral Flo and top up with water. Agitate gently but thoroughly before spraying commences and avoid over-vigorous agitation at any stage.

Rovral Flo is unstable in alkaline solutions: if alkaline water is used to prepare spray solutions use substitute Xchange® to reduce the pH of the final solution to below pH 7.

Rovral Flo is compatible with most commonly used fungicides and insecticides, but do not mix with strongly alkaline materials. If unsure, check compatibility before use.

INSTRUCTIONS FOR FIELD USE

Rates recommended below are for high volume spraying to the point of run off. Rates should be increased for concentrate spraying. Do not use less than 150 ml Rovral Flo/100 litres (3 litres/hectare).

It is an offence for users of this product to cause residues exceeding the relevant MRL in the New Zealand (Maximum Residues Limits of Agricultural Compounds) Food Standards.

CONDITIONS OF SUPPLY

READ THESE CONDITIONS OF SUPPLY BEFORE BUYING AND USING. IF THESE CONDITIONS ARE UNACCEPTABLE, RETURN AT ONCE UNOPENED.

To the extent permitted by law Bayer does not accept any liability or responsibility whatsoever for any loss, damage or injury whether direct, indirect or consequential howsoever arising in connection with these goods, including without limitation their storage, application, handling or use. Bayer does not make any warranties (express or implied) with respect to fitness for purpose, merchantable quality or any other matter.

To the extent only that Bayer is prevented by law from contracting out of the Consumer Guarantees Act, then these conditions apply subject to that Act. The Act shall not apply where these goods are acquired for business purposes.